

NEW YORK

NON-NATIVE PLANT INVASIVENESS RANKING FORM

Scientific name: Dipsacus laciniatus L. USDA Plants Code: DILA4
 Common names: Cut-leaf teasel
 Native distribution: Temperate Eurasia
 Date assessed: October 22, 2009
 Assessors: Gerry Moore
 Reviewers: LIISMA SRC
 Date Approved: November, 4, 2009 Form version date: 25 September 2009

New York Invasiveness Rank: High (Relative Maximum Score 70.00-80.00)

Distribution and Invasiveness Rank (<i>Obtain from PRISM invasiveness ranking form</i>)		
Status of this species in each PRISM:	Current Distribution	PRISM Invasiveness Rank
1 Adirondack Park Invasive Program	Not Assessed	Not Assessed
2 Capital/Mohawk	Not Assessed	Not Assessed
3 Catskill Regional Invasive Species Partnership	Not Assessed	Not Assessed
4 Finger Lakes	Not Assessed	Not Assessed
5 Long Island Invasive Species Management Area	Not Present	Low
6 Lower Hudson	Not Assessed	Not Assessed
7 Saint Lawrence/Eastern Lake Ontario	Not Assessed	Not Assessed
8 Western New York	Not Assessed	Not Assessed

Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 (<u>30</u>)	20
2	Biological characteristic and dispersal ability	25 (<u>25</u>)	16
3	Ecological amplitude and distribution	25 (<u>25</u>)	25
4	Difficulty of control	10 (<u>10</u>)	7
	Outcome score	100 (<u>90</u>) ^b	68 ^a
	Relative maximum score †		75.56
	New York Invasiveness Rank §	High (Relative Maximum Score 70.00-80.00)	

* For questions answered “unknown” do not include point value in “Total Answered Points Possible.” If “Total Answered Points Possible” is less than 70.00 points, then the overall invasive rank should be listed as “Unknown.”

† Calculated as 100(a/b) to two decimal places.

§ Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00
 Not Assessable: not persistent in NY, or not found outside of cultivation.

A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

A1.1. Has this species been documented to persist without cultivation in NY? (reliable source; voucher not required)		
<input checked="" type="checkbox"/>	Yes – continue to A1.2	
<input type="checkbox"/>	No – continue to A2.1	
A1.2. In which PRISMs is it known (see inset map)?		
<input checked="" type="checkbox"/>	Adirondack Park Invasive Program	
<input checked="" type="checkbox"/>	Capital/Mohawk	
<input type="checkbox"/>	Catskill Regional Invasive Species Partnership	
<input checked="" type="checkbox"/>	Finger Lakes	
<input type="checkbox"/>	Long Island Invasive Species Management Area	
<input type="checkbox"/>	Lower Hudson	
<input type="checkbox"/>	Saint Lawrence/Eastern Lake Ontario	
<input checked="" type="checkbox"/>	Western New York	

NEW YORK

NON-NATIVE PLANT INVASIVENESS RANKING FORM

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2009; Weldy & Werier, 2009.

A2.1. What is the likelihood that this species will occur and persist outside of cultivation, given the climate in the following PRISMs? (obtain from PRISM invasiveness ranking form)

Not Assessed	Adirondack Park Invasive Program
Not Assessed	Capital/Mohawk
Not Assessed	Catskill Regional Invasive Species Partnership
Not Assessed	Finger Lakes
Moderately Likely	Long Island Invasive Species Management Area
Not Assessed	Lower Hudson
Not Assessed	Saint Lawrence/Eastern Lake Ontario
Not Assessed	Western New York

Documentation:

Sources of information (e.g.: distribution models, literature, expert opinions):

Brooklyn Botanic Garden, 2009.

If the species does not occur and is not likely to occur in any of the PRISMs, then stop here as there is no need to assess the species. Rank is "Not Assessable."

A2.2. What is the current distribution of the species in each PRISM? (obtain rank from PRISM invasiveness ranking forms)

	Distribution
Adirondack Park Invasive Program	Not Assessed
Capital/Mohawk	Not Assessed
Catskill Regional Invasive Species Partnership	Not Assessed
Finger Lakes	Not Assessed
Long Island Invasive Species Management Area	Not Present
Lower Hudson	Not Assessed
Saint Lawrence/Eastern Lake Ontario	Not Assessed
Western New York	Not Assessed

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2009; Weldy & Werier, 2009.

A2.3. Describe the potential or known suitable habitats within New York. Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

<p>Aquatic Habitats</p> <p><input type="checkbox"/> Salt/brackish waters</p> <p><input type="checkbox"/> Freshwater tidal</p> <p><input type="checkbox"/> Rivers/streams</p> <p><input type="checkbox"/> Natural lakes and ponds</p> <p><input type="checkbox"/> Vernal pools</p> <p><input type="checkbox"/> Reservoirs/impoundments*</p>	<p>Wetland Habitats</p> <p><input type="checkbox"/> Salt/brackish marshes</p> <p><input checked="" type="checkbox"/> Freshwater marshes</p> <p><input checked="" type="checkbox"/> Peatlands</p> <p><input checked="" type="checkbox"/> Shrub swamps</p> <p><input type="checkbox"/> Forested wetlands/riparian</p> <p><input checked="" type="checkbox"/> Ditches*</p> <p><input type="checkbox"/> Beaches and/or coastal dunes</p>	<p>Upland Habitats</p> <p><input type="checkbox"/> Cultivated*</p> <p><input checked="" type="checkbox"/> Grasslands/old fields</p> <p><input type="checkbox"/> Shrublands</p> <p><input checked="" type="checkbox"/> Forests/woodlands</p> <p><input type="checkbox"/> Alpine</p> <p><input checked="" type="checkbox"/> Roadsides*</p>
---	---	---

Other potential or known suitable habitats within New York:

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2009.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

B. INVASIVENESS RANKING

Questions apply to areas similar in climate and habitats to New York unless specified otherwise.

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score

U

Documentation:
 Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)
 While the plant has been known in the U.S. since the 1800s, specific studies on its impacts to ecosystem processes and system wide parameters are not known
 Sources of information:
 Grauver, 2006; author's pers. comm.

1.2. Impact on Natural Community Structure

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score

7

Documentation:
 Identify type of impact or alteration:
 Large stands can significantly increase the density of the herb layer, and also significantly increasing the height of the herb layer, D. laciniatus obtaining heights up to 3 m. Snyder & Kaufman (2004): "teasels significantly alter the structure of rare natural plant communities."
 Sources of information:
 Snyder & Kaufmann, 2004; Gravuer, 2006; ; author's pers. obs.

1.3. Impact on Natural Community Composition

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) 10

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

U. Unknown

Score

10

Documentation:

Identify type of impact or alteration:

Smaller stands simply reduce the number of native individuals for a species in an area, whereas larger, dense stands can significantly reduce numbers of plant species. Also, in New Jersey, the species has been reported to have invaded limestone fens and caused the reduction or extirpation of several rare plant species, including the globally rare globe flower (*Trollius laxus* subsp. *laxus*).

Sources of information:

Snyder & Kaufman, 2004; author's pers. obs

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades. Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- | | |
|---|----|
| A. Negligible perceived impact | 0 |
| B. Minor impact | 3 |
| C. Moderate impact | 7 |
| D. Severe impact on other species or species groups | 10 |
| U. Unknown | |

Score

3

Documentation:

Identify type of impact or alteration:

Plant is quite prickly; other impacts to other species or species groups not known.

Sources of information:

Author's pers. obs.

Total Possible	<table border="1" style="display: inline-table;"><tr><td style="width: 50px; text-align: center;">30</td></tr></table>	30
30		
Section One Total	<table border="1" style="display: inline-table;"><tr><td style="width: 50px; text-align: center;">20</td></tr></table>	20
20		

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction

- | | |
|--|---|
| A. No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). | 0 |
| B. Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) | 1 |
| C. Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) | 2 |
| D. Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.) | 4 |
| U. Unknown | |

Score

4

Documentation:

Describe key reproductive characteristics (including seeds per plant):

Large plants can produce over 2,000 seeds; viability is high -- often 80% or more.

Sources of information:

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

Grauver, 2006; author's pers. obs.

2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair, buoyant fruits, pappus for wind-dispersal)

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score

Documentation:

Identify dispersal mechanisms:

Most seeds fall near the parent plant. Occasional long distance dispersal by water and wind -- possibly assisted by highways that create wind corridors -- may occur despite specific adaptations.

Sources of information:

Glass, 1990; Smith, 2004; Grauver, 2006; Rector et al., 2006; author's pers. obs.

2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

Score

Documentation:

Identify dispersal mechanisms:

Readily dispersed by mowing equipment; also occasionally sold for cultivation and in dried flower displays.

Sources of information:

Gremaud & SMith, 2002; Snyder & Kaufman, 2004; Grauver, 2006; author's pers. obs.

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 3
- C. Possesses two or more characteristics that increase competitive advantage 6
- U. Unknown

Score

Documentation:

Evidence of competitive ability:

Perennial or biennial monocarp (i.e., dies after setting seed); no other characteristics known that increase competitive advantage.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Sources of information:
Grauver, 2006.

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
- B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
- U. Unknown

Score 2

Documentation:

Describe growth form:
Forms a very dense tall thickety layer above shorter vegetation.
Sources of information:
Snyder & Kaufman, 2004; Grauver, 2006; author's pers. obs.

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
- B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
- C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
- U. Unknown (No studies have been completed)

Score 3

Documentation:

Describe germination requirements:
Germinates in existing vegetation, including fescue turfs, in a wide range of conditions, usually in richer, limestone soils.
Sources of information:
Grauver, 2006; author's pers. obs.

2.7. Other species in the genus invasive in New York or elsewhere

- A. No 0
- B. Yes 3
- U. Unknown

Score 0

Documentation:

Species:
Dipsacus fullonum, Dipsacus sativum, Dipsacus sylvestris. None tracked as invasive here or elsewhere. D. laciniatus is not as common as the common teasel (D. sylvestris). Brooklyn Botanic Garden, 2009; Weldy & Werier, 2009; U.S.D.A. NRCS, 2009.

Total Possible 25
Section Two Total 16

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: "The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude")

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

- B. Large dense stands present in areas with numerous invasive species already present or disturbed landscapes 2
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score

Documentation:

Identify reason for selection, or evidence of weedy history:
Large stands can occur over 0.25 acres sometimes in areas lacking other invasives.
Sources of information:
Snyder & Kaufman, 2004; Gravuer, 2006; author's pers. obs.

3.2. Number of habitats the species may invade

- A. Not known to invade any natural habitats given at A2.3 0
- B. Known to occur in one natural habitat given at A2.3 1
- C. Known to occur in two natural habitats given at A2.3 2
- D. Known to occur in three natural habitat given at A2.3 4
- E. Known to occur in four or more natural habitats given at A2.3 6
- U. Unknown

Score

Documentation:

Identify type of habitats where it occurs:
See A2.3.
Sources of information:
Snyder & Kaufman, 2004; Grauver, 2006; Brooklyn Botanic Garden, 2009; author's pers. obs.

3.3. Role of disturbance in establishment

- A. Requires anthropogenic disturbances to establish. 0
- B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
- C. Can establish independent of any known natural or anthropogenic disturbances. 4
- U. Unknown

Score

Documentation:

Identify type of disturbance:
Usually found in disturbed areas, but also reported from undisturbed areas such as fens.
Sources of information:
Snyder & Kaufman, 2005; Grauver, 2006; author's pers. obs.

3.4. Climate in native range

- A. Native range does not include climates similar to New York 0
- B. Native range possibly includes climates similar to at least part of New York. 1
- C. Native range includes climates similar to those in New York 3
- U. Unknown

Score

Documentation:

Describe what part of the native range is similar in climate to New York:
Temperate Europe and Asia.
Sources of information:
Grauver, 2006; Brooklyn Botanic Garden, 2009.

3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope)

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces. and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score

Documentation:

Identify states and provinces invaded:

KY, IN, IL, IA, MA, MD, MI, MN, MO, NJ, NY, OH, PA, VA, WI, WV; Ont.

Sources of information: See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.

U.S.D.A. NRCS, 2009.

3.6. Current introduced distribution of the species in natural areas in the eight New York State PRISMs (Partnerships for Regional Invasive Species Management)

- A. Present in none of the PRISMs 0
- B. Present in 1 PRISM 1
- C. Present in 2 PRISMs 2
- D. Present in 3 PRISMs 3
- E. Present in more than 3 PRISMs or on the Federal noxious weed lists 4
- U. Unknown

Score

Documentation:

Describe distribution:

See A1.1.

Sources of information:

Brooklyn Botanic Garden, 2009; Weldy & Werier, 2009.

Total Possible
Section Three Total

4. DIFFICULTY OF CONTROL

4.1. Seed banks

- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
- B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
- C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
- U. Unknown

Score

Documentation:

Identify longevity of seed bank:

Seeds reported to remain viable in soil for up to two years; no evidence for 10 years.

Sources of information:

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Glass, 1990; Smith, 2004.

4.2. Vegetative regeneration

- | | | |
|----|---|---|
| A. | No regrowth following removal of aboveground growth | 0 |
| B. | Regrowth from ground-level meristems | 1 |
| C. | Regrowth from extensive underground system | 2 |
| D. | Any plant part is a viable propagule | 3 |
| U. | Unknown | |

Score 1

Documentation:

Describe vegetative response:

Regrowth from basal rosettes.

Sources of information:

Grauver, 2006; author's pers. obs.

4.3. Level of effort required

- | | | |
|----|---|---|
| A. | Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. | 0 |
| B. | Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft ²). | 2 |
| C. | Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). | 3 |
| D. | Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). | 4 |
| U. | Unknown | |

Score 4

Documentation:

Identify types of control methods and time-term required:

The following is from Gruver (2006): "The Invasive Plant Association of Wisconsin (IPAW) regards this species as relatively difficult to control (IPAW 2003). Mechanical control is recommended in natural areas. In small stands, rosettes can be dug up, although plants often resprout if the root is not completely removed and damage to the surrounding area can occur if plants are large. Stalks can also be cut once flowering has begun, but before seed set. Because seeds can develop on immature heads, however, the cut stalks need to be removed from the area. Also, cutting of flowering stems may need to be repeated for several years to achieve effective control. Mowing is not an effective control, and in fact often increases the size of patches (Parrish et al. 2005). If mechanical control is not feasible, foliar application of herbicides can be used. Because rosettes of this species are green in early spring and late fall when many native plants are dormant, herbicide control during these times will minimize damage to native species. Also, dicot-selective herbicides (e.g. Triclopyr) are effective, which reduces damage to native monocots. As with mechanical control, however, herbicide applications over several years are required to manage an established population. Periodic prescribed burning may be helpful in conjunction with mechanical and/or chemical control (Glass 1990, Weber 2003, Smith 2004, WIDNR 2004, Czarapata 2005). No biocontrol agents are currently in use, but these are being researched (Rector et al. 2006).

"Several years (up to 5-6) of treatment may be necessary to totally eradicate this species from a natural community, regardless of whether mechanical or chemical treatment is chosen (Glass 1990, Gremaud and Smith 2002, Smith 2004, WIDNR 2004, Czarapata 2005).

"If rosettes are dug up, damage to the surrounding area can occur if plants are large. If

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

flowering stems are cut, native species of similar height may also be cut in the process. If herbicides are used, non-target damage may occur, though this can be minimized by spraying during the dormant season and/or using a dicot-specific herbicide (Glass 1990, Weber 2003, Smith 2004, WIDNR 2004, Czarapata 2005)."

Sources of information:
Childs, 2003; Fellows, 2004.

Total Possible	10
Section Four Total	7

Total for 4 sections Possible	90
Total for 4 sections	68

C. STATUS OF CULTIVARS AND HYBRIDS:

At the present time (May 2008) there is no protocol or criteria for assessing the invasiveness of cultivars independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain, and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Some cultivars of the species known to be available:

References for species assessment:

Brooklyn Botanic Garden. 2008. AILANTHUS database. [Accessed on October 22, 2009].

Cooperrider, T.S. 1995. The Dicotyledoneae of Ohio. Ohio State University Press, Columbus.

Czarapata, E. J. 2005. Invasive Plants of the Upper Midwest. The University of Wisconsin Press. Madison, WI. 215 pp.

Deam, C.M. 1940. Flora of Indiana. Division Forestry, Dept. Conservation, Indianapolis, Indiana. 1236 pp.

Glass, W. 1990. Vegetation management guideline: Cut-leaved teasel (*Dipsacus laciniatus* L.), Common teasel (*Dipsacus sylvestris* Huds.). Vol. 1, No. 24. Illinois Nature Preserves Commission. <inhs.uiuc.edu/chf/outreach/VMG/teasel.html>. [Accessed October 22, 2009.]

Gravuer, K. 2006. *Dipsacus laciniatus*. U.S. Invasive Species Impact Rank (I-Rank). NatureServe Explorer. <www.natureserve.org>. [Accessed on October 20, 2009.]

Gremaud, G. and T. Smith. 2002. Teasel Alert! Common and cut-leaved teasels - two species - one BIG problem! Missouri Department of Conservation. <mdc.mo.gov/documents/nathis/invasive/teasel.pdf> [Accessed October 22, 2009.]

Hilty, J. 2006. Illinois wildflowers. <illinoiswildflowers.info> [Accessed October 22, 2009.]

NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM

Invasive Plants Association of Wisconsin (IPAW). 2003. IPAW working list of the invasive plants of Wisconsin: a call for comments and information. *Plants Out of Place*, Issue 4. <ipaw.org/newsletters/issue4.pdf> [Accessed October 22, 2009].

Jones, R. L. 2005. *Plant Life of Kentucky*. The University Press of Kentucky. 834 pp.

Musser, A. and J. Parrish. 2002. Differences in *Dipsacus laciniatus* seed dispersal along an interstate corridor versus a state natural area. Abstract of poster presented at 2002 Ecological Society of America meeting, Tuscon, Arizona. <abstracts.co.allenpress.com/pweb/esa2002/document/?ID=16990> [Accessed October 22, 2009].

Ohio Department of Natural Areas and Parks (OHDNAP). 2001. *Invasive Plants of Ohio: Common and cut-leaved teasel*. Fact Sheet 15. <dnr.state.oh.us/dnap/invasive/pdf/InvasiveFactSheet15.pdf> [Accessed October 22, 2009].

Parrish, J., A. Oliver, R. Wiedenmann, S. Post, C. Helm, and M. Timpe. 2005. Effects of mowing on seed dispersal and patch growth of cut leafed teasel (*Dipsacus laciniatus*). Abstract of poster presented at 2005 Ecological Society of America meeting, Montreal, Canada. <abstracts.co.allenpress.com/pweb/esa2005/document/?ID=51710> [Accessed October 22, 2009.]

Peck, M.E. 1961. *A manual of the higher plants of Oregon*. 2nd edition. Binsford & Mort, Portland, Oregon. 936 pp.

Rector, B. G., V. Harizanova, R. Sforza, T. Widmer, and R. N. Wiedemann. 2006. Prospects for biological control of teasels, *Dipsacus* spp., a new target in the United States. *Biological Control* 36: 1-14.

Rhoads, A.F. and T.A. Block. 2000. *The Plants of Pennsylvania: An Illustrated Manual*. University of Pennsylvania Press: Philadelphia, Pennsylvania. 1061 pp.

Rice, P.M. 2006. *Invaders Database System*. Division of Biological Sciences, University of Montana, Missoula. <invader.dbs.umt.edu>. [Accessed October 22, 2009.]

Rydberg, Per Axel. 1932. *Flora of the Prairies and Plains of Central North America*. The New York Botanical Garden, New York, New York. 969 p.

Scoggan, H.J. 1978-1979. *The flora of Canada: Parts 1-4*. National Museums Canada, Ottawa. 1711 pp.

Seymour, F.C. 1989. *The flora of New England. A manual for the identification of all vascular plants including ferns and their allies growing without cultivation in New England*. Boston Museum Science, Boston. 611 pp.

Smith, T. E. 2004. *Missouri vegetation management manual*. Missouri Department of Conservation, Jefferson City. <mdc.mo.gov/nathis/exotic/vegman> [Accessed October 22, 2009.]

Steyermark, J.A. 1963. *Flora of Missouri*. Iowa State Univ. Press, Ames. 1728 pp.

Strausbaugh, P.D., and E.L. Core. 1978. *Flora of West Virginia*. Seneca Books, Inc., Grantsville, WV. 1079 pp.

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

- Swearingen, J. 2006. Alien plant invaders of natural areas. Plant Conservation Alliance, Alien Plant Working Group. <nps.gov/plants/alien/list> [Accessed on October 22, 2009.]
- Snyder, D. and S.R. Kaufman. 2004. An overview of nonindigenous plant species in New Jersey. New Jersey Department of Environmental Protection, Division of Parks and Forestry, Office of Natural Lands Management, Natural Heritage Program, Trenton, New Jersey. 107 pp.
- Tenaglia, D. 2006. The Missouri Flora Website. <missouriplants.com> [Accessed October 22, 2009.]
- The Nature Conservancy. 2001. Map: TNC Ecoregions of the United States. Modification of Bailey Ecoregions.
- United States Department of Agriculture, ARS, National Genetic Resources Program. 2005. December 9 last update. Germplasm Resources Information Network (GRIN) Online Database. National Germplasm Resources Laboratory, Beltsville, Maryland. <ars-grin.gov2/cgi-bin/npgs/html/index.pl>. [Accessed October 22, 2009.]
- United States Department of Agriculture, National Resources Conservation Service. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana. [Accessed on October 22, 2009.]
- Voss, E.G. 1996. Michigan Flora. Part III. Dicots (Pyrolaceae-Compositae). Cranbrook Institute of Science Bulletin 61 and Univ. Michigan Herbarium. Ann Arbor, Michigan. 622 pp.
- Weber, E. 2003. Invasive plant species of the world: a reference guide to environmental weeds. CABI Publishing, Cambridge, Massachusetts. 548 pp.
- Weber, W.A., and R.C. Wittmann. 1996. Colorado flora: Eastern slope. Revised edition. Univ. Press of Colorado, Niwot, Colorado. 524 pp.
- Weldy, T. and D. Werier. 2005. New York Flora Atlas. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. [Accessed on July 31, 2009.]
- Whiting, D. 2006. Dave's Garden: PlantFiles. <davesgarden.com/pf/> [Accessed October 22, 2009.]
- Wisconsin Department of Natural Resources (WDNR). 2004, September 3 last update. Cut-leaved teasel (*Dipsacus laciniatus*). Invasive species fact sheet. <dnr.state.wi.us/invasives/fact/teasel_cut.htm> [Accessed October 22, 2009.]

Citation: This NY ranking form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

Acknowledgments: The NY form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form. Original members of the LIISMA SRC included representatives of the Brooklyn Botanic Garden; The Nature Conservancy; New York Natural Heritage Program, New York Sea Grant; New York State Office of Parks, Recreation and Historic Preservation; National Park Service; Brookhaven National Laboratory; New York State Department of Environmental Conservation

NEW YORK

NON-NATIVE PLANT INVASIVENESS RANKING FORM

Region 1; Cornell Cooperative Extension of Suffolk/Nassau Counties; Long Island Nursery and Landscape Association; Long Island Farm Bureau; SUNY Farmingdale Ornamental Horticulture Department; Queens College Biology Department; Long Island Botanical Society; Long Island Weed Information Management System database manager; Suffolk County Department of Parks, Recreation and Conservation; Nassau County Department of Parks, Recreation and Museums; Suffolk County Soil & Water Conservation District.

References for ranking form:

- Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.
- Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).
- Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>
- Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49
- Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.
- Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.