

**PRISM**  
 (New York Partnerships for Regional Invasive Species Management)  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

PRISM: Long Island Invasive Species Management Area

Scientific name: Lonicera japonica Thunberg      USDA Plants Code: LOJA  
 Common names: Japanese Honeysuckle  
 Native Distribution: Eastern Asia  
 Date Assessed: 26 February 2008; edited 4 September 2009  
 PRISM Assessors: Steve Glenn  
 PRISM Reviewers: LIISMA Scientific Review Committee  
 Date Approved: May 14, 2008      Form version date: 13 April 2009  
 New York Relative Maximum score: 83.51      Date NY assessment approved: May 14, 2008  
 New York State Invasive Rank: Very High

**SUMMARY OF PRISM RANKING RESULTS:**

**Distribution:** Widespread  
**Estimated number of infested sites:** 100+  
**PRISM Invasiveness Rank<sup>§</sup>:** Very High



**A. DISTRIBUTION AND ABUNDANCE (KNOWN/POTENTIAL):**

1. What is the species distribution and abundance in the PRISM?

- |  |             |
|--|-------------|
| A. Not present   | Not Present |
| B. Occurs in three or fewer natural areas (locations that are at least ¼ mile apart) with no infested area* >1 acre or containing >100 individuals | Restricted  |
| C. Present in 4–10 natural areas, or with one occupied location >1 acre or containing >100 individuals   | Common      |
| D. Present in >10 minimally managed areas  | Widespread  |
| U. Unknown   | Unknown     |

Answer: Widespread

Describe distribution:  
 Widespread throuout PRISM.  
 Sources of information:  
 Brooklyn Botanic Garden. 2008.

<sup>§</sup>Not Assessable: not persistent in the PRISM, or not found outside of cultivation.

\*Definition of “infested area” is the “...actual or percentage of land occupied by [canopy cover of] weed plants”  
 NAWMA (North American Weed Management Association) 2002. North American Invasive Plant Mapping Standards (see <http://www.nawma.org/>).

**PRISM**  
(New York Partnerships for Regional Invasive Species Management)  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

2. What is the likelihood the species will occur (if not yet present) or expand its distribution and abundance (if already present) in the PRISM?

Answer:

Documentation (e.g.: history of establishment in PRISM, suitability of habitats and climate, distribution models, literature, expert opinions):  
Current widespread distribution.  
Sources of information:  
Brooklyn Botanic Garden. 2008.

**B. INVASIVENESS RANK IN THE PRISM:**

Is the species distribution Widespread or Common?

Yes: Go to column A in table below.

No: What is the likelihood of species occurrence or expansion? Answer:

- Very Likely:                      Use column A below
- Moderately likely:            Use column B below
- Unlikely:                         Use column C below
- Zero likelihood                 Invasive potential Insignificant
- Unknown                         Invasive potential Unknown
- Not assessed                     Invasive potential not assessed

Assign a PRISM invasiveness rank to the species based on its New York Relative Maximum Score, using the designated column in the table below.

New York Relative Maximum Score	New York Invasiveness Rank	A	B	C
> 80.00	Very High	VH	H	M
70.00–80.00	High	H	M	L
50.00–69.99	Moderate	M	L	Ins
40.00–49.99	Low	L	Ins	Ins
<40.00	Insignificant	Ins	Ins	Ins

**Column used: A (Insert PRISM Invasiveness Rank on page 1)**

**References for species assessment:**

Brooklyn Botanic Garden. 2008. AILANTHUS databse. (accessed 26 February 2008).

Fowler, S. P. & K. C. Larson. 2004. Seed germination and seedling recruitment of Japanese honeysuckle in a central Arkansas natural area. *Natural Areas J.* 24: 49-53.

Huebner, C. D. 2003. Vulnerability of oak-dominated forests in West Virginia to invasive exotic plants: Temporal and spatial patterns of nine exotic species using herbarium records and land classification data. *Castanea* 68:1-14.

Larson, B. M. H. et al. 2007. The biology of Canadian weeds. 135. *Lonicera japonica* Thunb. *Canad. J. Pl. Sci.* 87: 423-438.

Larson, K. C. 2000. Circumnutation behavior of an exotic honeysuckle vine and its native congener: Influence on clonal mobility. *Amer. J. Bot.* 87: 533-538.

**PRISM**  
(New York Partnerships for Regional Invasive Species Management)  
**NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- Li, Q. et al. 2007. Leaf epidermal characters of *Lonicera japonica* and *Lonicera confusa* and their ecology adaptation. *J. Forestry Res. (Harbin)* 18: 103-108.
- Hidayati, S. N. et al. 2000. Dormancy-breaking and germination requirements of seeds of four *Lonicera* species (Caprifoliaceae) with underdeveloped spatulate embryos. *Seed Sci. Res.* 10: 459-469.
- Honu, Y. A. K. & D. J. Gibson. 2006. Microhabitat factors and the distribution of exotic species across forest edges in temperate deciduous forest of southern Illinois, USA. *J. Torrey Bot. Soc.* 133: 255-266.
- New York Flora Association. 2008. New York Flora Atlas. <<http://atlas.nyflora.org/>> (accessed 26 February 2008).
- Morrow, J. L. et al. 2001. Habitat selection and habitat use by the bog turtle (*Clemmys muhlenbergii*) in Maryland. *J. Herpetology* 35: 545-552.
- Nuzzio, V. 1997. Element stewardship abstract for *Lonicera japonica*. The Nature Conservancy, Arlington, VA.
- Rehehr, D. L. & D. R. Frey. 1988. Selective control of Japanese Honeysuckle (*Lonicera japonica*). *Weed Tech.* 139-143.
- Schierenbeck, K. A. & J. D. Marshall. 1993. Seasonal and diurnal patterns of photosynthetic gas exchange for *Lonicera sempervirens* and *L. japonica* (Caprifoliaceae). *Amer. J. Bot.* 80: 1292-1299.
- Skulman, B. W. et al. 2004. Evidence for allelopathic interference of Japanese honeysuckle (*Lonicera japonica*) to loblolly and shortleaf pine regeneration. *Weed Science* 52: 433-439.
- Suthers, H. B. et al. 2000. Use of successional habitat and fruit resources by songbirds during autumn migration in central New Jersey. *Wilson Bull.* 112: 249-260.
- United States Department of Agriculture. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, LA <<http://plants.usda.gov>> (accessed 26 February 2008)
- Wang, F. & W. C. Ma. 2004. Comparative studies on light utilization characteristics and shade tolerance of 7 climbing shrub species. *Forest Research* 17: 305-309.
- Webb, S. L. et al. 2000. The myth of the resilient forest: Case study of the invasive Norway maple (*Acer platanoides*). *Rhodora* 102(911): 332-354. [*L. japonica* also discussed]
- White, D. W. & Stiles, E. W. 1992. Bird dispersal of fruits of species introduced into eastern North America *Canad. J. Bot.* 70: 1689-1696.
- Williams, C. E. 1999. Fruits of alien shrubs and deer mice: A test of the persistent fruit defense hypothesis. *J. Penn. Acad. Sci.* 73: 33-37.
- Wright, D. E. & J. F. Pagels. 1977. Climbing activity in the hispid cotton rat, *Sigmodon hispidus*, and the eastern meadow mole, *Microtus pennsylvanicus*. *Chesapeake Sci.* 18: 87-89.
- Yukonis, K. A. & S. J. Meiners. 2004. Invasion impacts local species turnover in a successional system. *Ecology Letters* 7: 764-769.

**Citation:** This ranking form for regions within NYS may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

**Acknowledgments:** Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form.